

Survival by Degrees: 389 Species on the Brink

Background

Birds form part of healthy ecosystems, bring joy to people, and benefit local economies throughout the United States. In 2011, birdwatching-related industries drove \$41 billion in expenditures and \$107 billion in total industry output nationally. There are more than 1,188,000 total birders in Colorado alone [1]. Additionally, birds play critical roles in pollination, insect control, forest generation, seed dispersal, carrion scavenging, and many other ecosystem services we rely on.

However, the future of birds is at risk with alarming losses of biodiversity occurring worldwide. Global extinction rates are now 100 times higher than background rates [2]. Climate change exacerbates the global biodiversity crisis, with an anticipated rate of change 20 times faster in the next century than during the past two million years.

Audubon leads the way in conducting science to understand the vulnerability and threats to birds from climate change. Our science shows that stabilizing warming at a global average of 1.5°C (2.7°F), as recommended by the IPCC (Intergovernmental Panel on Climate Change) to reduce the global risk of climate change, would also reduce vulnerability and threats for many species of birds. To save birds we must address the underlying causes of climate change (*climate change mitigation*), and protect places that birds need now and will need in the future (*climate change adaptation*). Climate change mitigation means reducing or preventing the causes of climate change, such as greenhouse gas emissions. Climate change adaptation includes efforts to alter and adapt both our natural surroundings as well as our infrastructure to better withstand the threats of climate change.

Audubon's 2019 Report, *Survival by Degrees: 389 Bird Species on the Brink* [3], is a powerful look at how vulnerable birds are to climate change across North America based on a new, updated scientific analysis that leverages big data and incorporates the unique biology of each bird to determine its vulnerability. In this research, we related bird observations for 604 species with climate and habitat conditions at these locations and used modeling algorithms to capture the unique composition of each species's suitable range. We then mapped and compared the projected current and future ranges to estimate the projected range loss and gain under multiple future climate change scenarios. These projections were then used to assess how vulnerable each species was to climate change [4,5].



Figure 1. Lark Bunting. Photo: Evan Barrientos/Audubon

Future Climate and Habitat in Colorado

Across the state of Colorado, without substantial climate change mitigation (i.e., a 3°C/5.4°F global warming scenario), average temperatures during the warmest month are expected to increase approximately 6.1°C (11°F), and average temperatures during the coldest month are expected to increase approximately 4°C (7.2°F) from 2010 to the end of the century. Average annual precipitation is expected to decrease by 9.9 mm (0.4 in), ranging from a decrease of 68 mm (2.7 in) in the east to an increase of 268 mm (10.5 in) in the north of the state. In addition, evapotranspiration is expected to increase, resulting in an overall decrease of available moisture of 43% across the state [6].

The distribution of vegetation biomes, critical for plants and animals, are also projected to change under climate change scenarios [7]. By the end of the century under a 3°C (5.4°F) global warming scenario, approximately 42% of the state of Colorado will transition to a different biome. At present, the largest biome in the state is Grassland, covering 40% of the state. By the end of the century, Grassland will cover approximately 45% of the state.

All of these changes in climate and vegetation will alter plant and insect communities; influence availability of food, water, and shelter for birds; and will likely cause ecological disruption as species assemblages reshuffle. Over time, a complex suite of changes in climate and vegetation will inevitably affect Colorado's bird communities.

Climate Change Vulnerability

Climate change will negatively affect many birds in the state. Here, we assess vulnerability based on the amount of a species's range that may be gained or lost with climate change. We designate species that may lose much more range across North America than they have the potential to gain as *climate vulnerable*. In Colorado, 123 out of 241 species are climate vulnerable in summer under the 3°C scenario, meaning they stand to lose more of their North American summer range than they would gain under a warming climate. Reducing emissions to 1.5°C reduces the number of vulnerable species to 84. Impacts are somewhat lessened in winter, with 39 out of 152 species vulnerable

under 3°C of warming and 21 species vulnerable if we reduce warming to 1.5°C.

Each bird was grouped by its primary habitat (see Table 2 for groupings), and these groups are not equally affected. In Colorado, the habitat groups with the most species vulnerable to the impacts of ongoing and future climate change are western forest (42 species) and boreal forest (18 species) in summer (Figure 2). In winter, western forest (15 species) and boreal forest (9 species) groups have the most vulnerable species.




Figure 2. Number of species by their vulnerability to climate change in each habitat group under a global 3°C warming scenario. The species in each group are ones that currently live in the state, though vulnerability is assessed across the species's full North American range to better account for range-wide changes. Red and orange indicate number of vulnerable (high and moderate) species, and yellow and blue indicate non-vulnerable (low and neutral) species.

Climate-Related Threats

In addition to changes in climate across North America, we assessed the potential impacts of other forecasted threats related to climate change, including sea level rise, land use change, and extreme weather events, such as extreme spring heat, spring drought, fire weather, heavy rain, and false springs within the lower 48 states [8]. These threats are relevant to both birds and the places they need, but were only available for the lower 48 states, and were analyzed separately from vulnerability. This analysis provides information on how each location and the birds that occur there may be exposed to these specific, climate-related threats (Figure 3) beyond their range-wide vulnerability described above.

Here we summarize threats occurring within the state. Seven climate-related threats will affect portions of Colorado (Table 1). The threat affecting both the greatest area and number of species in the state is extreme spring heat.

In Colorado, species that are most threatened by a combination of climate change and additional climate-related threats under 3°C of warming include Greater Sage-Grouse, Dusky Grouse, Broad-tailed Hummingbird, Mountain Plover, American Dipper, Lark Bunting, Mountain Chickadee, Black Rosy-Finch, and Clark's Nutcracker. For information on threats for individual species in Colorado, see Table 2.

Climate-Related Threats (Cont.)











Figure 3. The number and distribution of overlapping climate-related threats under future global change scenarios of 1.5°C and 3°C. For detailed information on threats for each location in the state, refer to our online interactive tool at climate.audubon.org.

Table 1. Climate-related threats that Colorado is expected to experience under the warming scenarios 1.5°C (2.7°F) and 3°C (5.4°F), and the projected area and number of species affected. We report the projected amount(s) of global sea level rise associated with each scenario [8]. Threats and scenarios were omitted if no species were affected in that scenario.

Threat	Scenario	Area Affected (acres)	Summer (Vulnerable) Species Affected	Winter (Vulnerable) Species Affected
 Urbanization	1.5°C	940,472	1 (0)	
	3°C	2,885,989	7 (2)	7 (1)
 Cropland Expansion	1.5°C	2,680,348	3 (2)	4 (0)
	3°C			
 Extreme Spring Heat	1.5°C	33,991,764	128 (58)	50 (14)
	3°C	66,660,527	234 (93)	189 (41)
 Fire Weather	1.5°C	11,418,475	8 (1)	4 (0)
	3°C	66,207,831	234 (93)	189 (41)
 Spring Droughts	3°C	4,918,625	1 (0)	
 Heavy Rain	1.5°C	5,705,902	3 (3)	1 (0)
	3°C	6,776,360	5 (2)	3 (2)
 False Springs	3°C	6,808,731	7 (0)	15 (2)

We also mapped risk, areas of high conservation value for birds that are exposed to climate change-related threats. For any one location, risk is the product of the number of overlapping climate change-related threats, the total number of bird species that occur under future climate, and

the number of species with range-wide vulnerability under future climate. Risk is greater across Colorado in summer relative to winter, and mitigating warming from 3°C to 1.5°C would more than halve the average risk of climate change-related threats to birds across the state.

Conclusions and Caveats

Birds are early responders to climate change and can be important indicators of large-scale ongoing and future ecological change. We found that 51% of Colorado's 270 bird species are vulnerable to climate change across seasons. A rapidly changing climate could lead to population declines and local extinctions if species are not able to adapt. In addition, the reshuffling of bird communities at a continental scale will bring together species that previously lived in isolation, leading to novel, unpredictable interactions. Disruptions in food and nesting resources further compound vulnerabilities to climate change.

Although we project range gains offsetting loss for some species, especially in winter, it is unknown whether birds will establish populations in these new locations because of other factors not assessed here. On top of this, the added stressors of extreme weather events and other climate change-related threats will make establishment and persistence of populations difficult in the coming decades.

Call to Action

We know what to do.

The scientific consensus is clear. We must reduce greenhouse gas emissions at an urgent speed and on a wide scale from every sector of the economy to achieve a more favorable future for birds and people. There is no single perfect solution, but we can make a series of changes that lead to large-scale, systemic adjustments to achieve the required reductions.

Addressing the underlying causes of climate change.

Audubon is pursuing policies that together can drive down emissions at the scale and speed we need. For instance, we can invest in 100% clean energy, energy efficiency, and clean transportation policies that will dramatically reduce carbon emissions from the U.S. and world economies. We can adapt, improve, and innovate. We can power our cars, homes, cities, factories, farms, communities, and economy with clean energy—without contributing to climate change. We are working to implement policies and conservation practices that offset what we cannot eliminate, such as planting forests and testing new technologies to capture (i.e., sequester) carbon through industrial processes and permanently store it underground. We can do all of this in ways that spur innovation, create good jobs, promote homegrown industries, and build our economy for a smarter future.

Protecting the places birds need.

We can also pursue policies and conservation practices that help us avoid some of the worst effects of climate change

While these studies did not assess the effects of climate change on people, we know that the fate of humans and birds are deeply connected. Climate change is currently and will continue to cause harm to people too, who face threats like extreme weather, loss of coastal areas and changing economic patterns, to name a few. Climate change will cause disproportionate harm to vulnerable communities, including children, the elderly, the sick, and the poor, who may have fewer resources available to move or otherwise protect themselves from these threats. If we drastically reduce carbon emissions, we help people and birds alike.

This is the most comprehensive assessment of climate change vulnerability of birds in North America to date, but even this assessment may reasonably be considered conservative because the pace of change is exceeding the scenarios considered in this study. Our work concludes that climate change will have multiple, compounding effects on birds and will likely amplify biodiversity loss, unless actions are taken to lessen its effects.

by building more resilient infrastructure—meaning our cities, roads, and other structures—or even ranches, parks, floodplains, forests, and wetlands that can serve as good wildlife habitat and simultaneously protect our communities from extreme weather.

Audubon has identified the best opportunities to increase the resilience of coastal wetlands in key places that can serve as the first line of defense against the threat of sea level rise. We work to ensure key landscapes that are critical for birds have clean and reliable sources of water, now and in the future, and we advocate for conservation-minded management of working and urban landscapes that can help birds adapt to the changing climate.

We still have time.

We can avert and limit dangerous warming and its worst effects if we act quickly. Science tells us that in order to limit warming to a rise of 1.5°C (2.7°F), we must reduce greenhouse gas emissions 45% below 2010 levels by 2030 and reach net-zero carbon emissions by 2050.

We must act now.

We are on a dangerous path, but we have the power to chart a better one. Still, change will come only if we demand action from the public officials who represent us and the businesses we support.

We ask you to join us.

Be part of the solution. We can do this, together.

How You Can Help in Colorado

We still have time.

Colorado still holds vast natural landscapes, from montane forests to flat grasslands and shrublands, and the rivers that connect them. Continuing to protect these remaining habitats and restore them where appropriate will help birds weather the impacts of climate change.

We must act now.

Fortunately, there are things everyone can do to fight climate change. Audubon can connect you to solutions making a difference across Colorado. Create urban refuges for birds by planting a bird-friendly garden. Keep carbon sequestered in grasslands by supporting ranches certified by Audubon’s Conservation Ranching Initiative. Advocate for policies that support healthy rivers and sagebrush steppe. Learn more at rockies.audubon.org.

More Information

This project was conducted by the National Audubon Society. For more information, including details on the methods, please see the project website (climate.audubon.org) and the scientific publications [5,8].

References

1. US Fish & Wildlife Service. 2013. Birding in the United States: A Demographic and Economic Analysis. Addendum to the 2011 National Survey of Fishing, Hunting, and Wildlife-Associated Recreation.

2. Ceballos, G. et al. 2015. Accelerated modern human-induced species losses: Entering the sixth mass extinction. *Science Advances* 1:e1400253. doi:10.1126/sciadv.1400253.

3. Wilsey, C. et al. 2019. Survival By Degrees: 389 Bird Species on the Brink. National Audubon Society: New York.

4. Wilsey, C. et al. 2019. Climate policy action needed to reduce vulnerability of conservation-reliant grassland birds in North America. *Conservation Science and Practice* e21; DOI: 10.1111/csp2.21.

5. Bateman, B. et al. 2019. North American birds require mitigation and adaptation to reduce vulnerability to climate change. In review.

6. AdaptWest Project. 2015. Gridded current and projected climate data for North America at 1km resolution, interpolated using the ClimateNA v5.10 software (T. Wang et al., 2015). Available at adaptwest.databasin.org.

7. Rehfeldt, G.E. et al. 2012. North American vegetation model for land-use planning in a changing climate: a solution to large classification problems.

8. Bateman, B. et al. 2019. Risk to North American birds from climate change-related threats. In review.

9. IPCC (Intergovernmental Panel on Climate Change). 2019. IPCC Special Report on the Ocean and Cryosphere in a Changing Climate.

Contact

Brooke Bateman, PhD
Senior Climate Scientist, National Audubon Society
climatescience@audubon.org

Species Projections





















































Table 2. Climate suitability projections in summer and winter under the 3°C warming scenario for birds in Colorado. Each bird is associated with the *Habitat Group* representing its primary habitat (see classification key below). *Range-wide Vulnerability* is the vulnerability of each species, across its full North American range under 3°C of global warming, based on long-term climate and vegetation change. High and moderately vulnerable species are considered vulnerable to climate change, whereas low and neutral species are considered not vulnerable. In *State Trends*, we show the top two trends in climate and habitat suitability for select birds in Colorado, with colors reflecting the trend according to the legend below and percentages reflecting the percent of the state’s area in which each trend will occur. The total percentage reflects the area of the state that the species currently occupies and is projected to occupy in the future. Potential colonization indicates that climate and habitat are projected to become suitable for the species, whereas potential extirpation indicates that climate and habitat are suitable today but projected to become unsuitable. *State Threats* shows the additional climate-related threats each species might face, indicated by icons as in Table 1. Threats shown here were assessed within each state for species under either 1.5°C or 3°C warming (i.e., species that will be completely extirpated from the state do not have threats shown). Omitted species are either not present in the state during that season or not modeled due to data deficiency. These lists may have been further reduced by local experts. For a full list of species modeled in Colorado, see the project website (climate.audubon.org).
















































Habitat classifications:



























































F-B = Boreal Forests
 F-E = Eastern Forests
 F-W = Western Forests
 F-S = Subtropical Forests
 A = Arctic
 D = Aridlands
 G = Grasslands
 M = Marshlands
 C = Coastal
 W = Waterbirds
 Gen = Generalists

Trend classifications:






 Potential extirpation
 Worsening
 Stable
 Improving
 Potential colonization





Species	Habitat Group	Season	Range-wide Vulnerability	State Trends	State Threats
Snow Goose	W	Winter	Low		 
Ross's Goose	W	Winter	Low		 
Cackling Goose	M	Winter	Moderate		 
Canada Goose	W	Summer	Moderate		 
	W	Winter	Neutral		 
Wood Duck	W	Summer	Low		 
	W	Winter	Neutral		 
Blue-winged Teal	M	Summer	Low		 
Cinnamon Teal	M	Summer	Moderate		 
	M	Winter	Neutral		 
Northern Shoveler	M	Summer	Low		 
	M	Winter	Neutral		 
Gadwall	M	Summer	Moderate		 
	M	Winter	Neutral		 
American Wigeon	M	Summer	Moderate		
	M	Winter	Neutral		 
Mallard	W	Summer	Low		 
	W	Winter	Neutral		 




















































Species	Habitat Group	Season	Range-wide Vulnerability	State Trends		State Threats	
Northern Pintail	M	Summer	Moderate	22%	7%		
	M	Winter	Neutral	37%	13%		
Green-winged Teal	M	Summer	Moderate	7%	6%		
	M	Winter	Neutral	31%	27%		
Canvasback	M	Winter	Neutral	29%	31%		
Redhead	M	Summer	Neutral	14%	8%		
	M	Winter	Low	18%	19%		
Ring-necked Duck	W	Summer	Moderate	16%	14%		
	W	Winter	Neutral	23%	22%		
Greater Scaup	W	Winter	Neutral	3%			
Lesser Scaup	W	Summer	High	9%	<1%		
	W	Winter	Neutral	23%	27%		
Bufflehead	W	Summer	High	4%			
	W	Winter	Low	10%	7%		
Common Goldeneye	W	Summer	High	5%	1%		
	W	Winter	Neutral	78%	12%		
Barrow's Goldeneye	W	Summer	High	3%	<1%		
Hooded Merganser	W	Summer	Low	2%			 
	W	Winter	Neutral	12%	5%		
Common Merganser	W	Summer	Moderate	26%	13%		
	W	Winter	Low	56%	30%		
Ruddy Duck	M	Summer	Low	24%	9%		
	M	Winter	Neutral	1%	16%		
Northern Bobwhite	G	Summer	Neutral	3%	7%		
	G	Winter	Neutral	<1%	6%		
Scaled Quail	D	Summer	Moderate	15%	31%		
	D	Winter	Moderate	7%	27%		
Ruffed Grouse	F-B	Summer	Moderate	7%			




















































Species	Habitat Group	Season	Range-wide Vulnerability	State Trends	State Threats		
Greater Sage-Grouse	F-B	Winter	Moderate	5%			
	D	Summer	High	10% 1%			
	D	Winter	High	15% 2%			
Dusky Grouse	F-W	Summer	High	20% 9%			
	F-W	Winter	High	24% 6%			
Sharp-tailed Grouse	G	Summer	Low	10%			
	G	Winter	Moderate	18%			
Greater Prairie-Chicken	G	Summer	Neutral	11% 8%			
	G	Winter	Neutral	8% 3%			
Lesser Prairie-Chicken	G	Summer	Moderate	5% 24%			
	G	Winter	Moderate	3% 23%			
Wild Turkey	Gen	Summer	Neutral	11% 20%			
	Gen	Winter	Neutral	20% 28%			
Pied-billed Grebe	M	Summer	Neutral	1% 5%			
	M	Winter	Neutral	1% 7%			
Horned Grebe	M	Winter	Neutral	1% 1%			
Eared Grebe	M	Summer	High	5% 6%			
	M	Winter	Neutral	1% 4%			
Western Grebe	M	Summer	Low	4% 11%			
	M	Winter	Low	1%			
Clark's Grebe	M	Summer	Low	1% 5%			
Inca Dove	D	Summer	Neutral	5%			
	D	Winter	Neutral	2%			
White-winged Dove	D	Summer	Neutral	4%			
Mourning Dove	Gen	Summer	Neutral	58% 27%			
	Gen	Winter	Neutral	41%			
Greater Roadrunner	D	Summer	Neutral	1% 18%			
	D	Winter	Neutral	<1% 15%			















Species	Habitat Group	Season	Range-wide Vulnerability	State Trends	State Threats		
Lesser Nighthawk	D	Summer	Neutral	<div><div></div>13%</div>			
Common Nighthawk	Gen	Summer	Neutral	<div><div></div>40%<div></div>50%</div>			
Common Poorwill	D	Summer	Neutral	<div><div></div>25%<div></div>33%</div>			
Black Swift	F-W	Summer	Moderate	<div><div></div>2%<div></div><1%</div>			
Chimney Swift	F-E	Summer	Neutral	<div><div></div>3%</div>			
White-throated Swift	D	Summer	Low	<div><div></div>6%<div></div>12%</div>			
Ruby-throated Hummingbird	F-E	Summer	Neutral	<div><div></div>3%</div>			
Black-chinned Hummingbird	D	Summer	Neutral	<div><div></div>21%<div></div>59%</div>			
Broad-tailed Hummingbird	F-W	Summer	High	<div><div></div>19%<div></div>17%</div>			
Calliope Hummingbird	F-W	Summer	High	<div><div></div>6%</div>			
Virginia Rail	M	Summer	Moderate	<div><div></div>17%<div></div>4%</div>			
	M	Winter	Low	<div><div></div>5%<div></div>16%</div>			
Sora	M	Summer	Moderate	<div><div></div>10%<div></div>1%</div>			
American Coot	M	Summer	Neutral	<div><div></div>5%<div></div>17%</div>			
	M	Winter	Neutral	<div><div></div>12%<div></div>39%</div>			
Black Rail	M	Summer	Moderate	<div><div></div>3%<div></div><1%</div>			
Sandhill Crane	M	Summer	Moderate	<div><div></div>13%<div></div>1%</div>			
	M	Winter	Low	<div><div></div>5%<div></div>18%</div>			
Black-necked Stilt	M	Summer	Neutral	<div><div></div>3%<div></div>23%</div>			
American Avocet	M	Summer	Neutral	<div><div></div>10%<div></div>14%</div>			
	M	Winter	Neutral	<div><div></div>1%</div>			
Snowy Plover	C	Summer	Neutral	<div><div></div>2%<div></div>30%</div>			
Killdeer	W	Summer	Neutral	<div><div></div>57%<div></div>6%</div>			
	W	Winter	Neutral	<div><div></div>9%<div></div>57%</div>			
Mountain Plover	G	Summer	High	<div><div></div>35%<div></div>6%</div>			
Upland Sandpiper	G	Summer	Neutral	<div><div></div>8%<div></div>11%</div>			
Long-billed Curlew	G	Summer	High	<div><div></div>42%<div></div>6%</div>			

Species	Habitat Group	Season	Range-wide Vulnerability	State Trends	State Threats		
Marbled Godwit	M	Summer	Moderate	<div><div><1%</div><div><1%</div></div>			
Wilson's Snipe	M	Summer	Moderate	<div><div>9%</div><div>1%</div></div>			
	M	Winter	Neutral	<div><div>18%</div><div>42%</div></div>			
Wilson's Phalarope	M	Summer	Low	<div><div>14%</div><div>3%</div></div>			
Spotted Sandpiper	W	Summer	Moderate	<div><div>26%</div><div>37%</div></div>			
Greater Yellowlegs	W	Winter	Neutral	<div><div>3%</div></div>			
Willet	W	Summer	Neutral	<div><div>3%</div><div>6%</div></div>			
Ring-billed Gull	W	Summer	Low	<div><div>5%</div><div>2%</div></div>			
	W	Winter	Neutral	<div><div>39%</div><div>20%</div></div>			
California Gull	W	Summer	Moderate	<div><div>16%</div><div>12%</div></div>			
	W	Winter	Low	<div><div>5%</div><div>18%</div></div>			
Herring Gull	W	Winter	Neutral	<div><div>43%</div><div>18%</div></div>			
Forster's Tern	M	Summer	Neutral	<div><div>4%</div></div>			
Neotropic Cormorant	C	Summer	Neutral	<div><div>2%</div></div>			
Double-crested Cormorant	W	Winter	Neutral	<div><div><1%</div><div>4%</div></div>			
American White Pelican	M	Summer	Low	<div><div>7%</div><div>5%</div></div>			
American Bittern	M	Summer	Low	<div><div>5%</div><div><1%</div></div>			
Great Blue Heron	W	Summer	Neutral	<div><div>62%</div><div>19%</div></div>			
	W	Winter	Neutral	<div><div>30%</div><div>34%</div></div>			
Great Egret	W	Summer	Neutral	<div><div>1%</div><div>2%</div></div>			
Snowy Egret	M	Summer	Neutral	<div><div><1%</div><div>2%</div></div>			
Cattle Egret	W	Summer	Neutral	<div><div>8%</div><div>27%</div></div>			
Black-crowned Night-Heron	M	Summer	Neutral	<div><div>28%</div><div>19%</div></div>			
	M	Winter	Neutral	<div><div><1%</div><div>8%</div></div>			
Yellow-crowned Night-Heron	M	Summer	Neutral	<div><div>43%</div></div>			
White-faced Ibis	M	Summer	Low	<div><div>9%</div><div>31%</div></div>			
Turkey Vulture	Gen	Summer	Neutral	<div><div>26%</div><div>31%</div></div>			

Species	Habitat Group	Season	Range-wide Vulnerability	State Trends		State Threats		
Osprey	W	Summer	Neutral	10% 2%				
Golden Eagle	Gen	Summer	Moderate	28% 45%				
	Gen	Winter	Moderate	3% 94%				
Mississippi Kite	F-E	Summer	Neutral	9% 35%				
Northern Harrier	M	Summer	Low	48% 5%				
	M	Winter	Neutral	47% 16%				
Sharp-shinned Hawk	F-W	Summer	Moderate	22% 20%				
	F-W	Winter	Neutral	60% 32%				
Cooper's Hawk	Gen	Summer	Neutral	34% 29%				
	Gen	Winter	Low	25% 46%				
Northern Goshawk	F-B	Summer	High	28% 16%				
	F-B	Winter	Low	23% 27%				
Bald Eagle	Gen	Summer	Low	44% 28%				
	Gen	Winter	Neutral	29% 58%				
Swainson's Hawk	G	Summer	Neutral	12% 46%				
Zone-tailed Hawk	F-W	Summer	Neutral	2%				
Red-tailed Hawk	Gen	Summer	Neutral	91% 8%				
	Gen	Winter	Neutral	54% 24%				
Rough-legged Hawk	A	Winter	Moderate	39% 27%				
Ferruginous Hawk	G	Summer	Moderate	30% 15%				
	G	Winter	Moderate	50% 11%				
Barn Owl	Gen	Summer	Neutral	24% 45%				
	Gen	Winter	Neutral	28% 43%				
Western Screech-Owl	F-W	Summer	Neutral	5% 30%				
	F-W	Winter	Neutral	4% 30%				
Eastern Screech-Owl	F-E	Summer	Neutral	22% 12%				
	F-E	Winter	Neutral	8%				
Great Horned Owl	Gen	Summer	Neutral	74% 11%				



















































Species	Habitat Group	Season	Range-wide Vulnerability	State Trends	State Threats		
	Gen	Winter	Neutral	<div><div></div><div>71%</div><div>12%</div></div>			
Snowy Owl	A	Winter	Low	<div><div></div><div>44%</div><div>2%</div></div>			
Northern Pygmy-Owl	F-W	Summer	High	<div><div></div><div>11%</div><div>9%</div></div>			
	F-W	Winter	High	<div><div></div><div>17%</div><div>12%</div></div>			
Burrowing Owl	G	Summer	Neutral	<div><div></div><div>47%</div><div>9%</div></div>			
	G	Winter	Neutral	<div><div></div><div>37%</div><div></div></div>			
Spotted Owl	F-W	Summer	High	<div><div></div><div>2%</div><div></div></div>			
Long-eared Owl	F-W	Summer	Low	<div><div></div><div>15%</div><div>66%</div></div>			
	F-W	Winter	Low	<div><div></div><div>53%</div><div>24%</div></div>			
Short-eared Owl	G	Summer	Moderate	<div><div></div><div>46%</div><div>8%</div></div>			
	G	Winter	Neutral	<div><div></div><div>17%</div><div>12%</div></div>			
Boreal Owl	F-B	Summer	High	<div><div></div><div>12%</div><div><1%</div></div>			
	F-B	Winter	High	<div><div></div><div>15%</div><div>1%</div></div>			
Northern Saw-whet Owl	F-B	Summer	Moderate	<div><div></div><div>29%</div><div>11%</div></div>			
	F-B	Winter	Low	<div><div></div><div>42%</div><div>25%</div></div>			
Belted Kingfisher	Gen	Summer	Neutral	<div><div></div><div>7%</div><div>14%</div></div>			
	Gen	Winter	Neutral	<div><div></div><div>47%</div><div>35%</div></div>			
Williamson's Sapsucker	F-W	Summer	High	<div><div></div><div>24%</div><div>12%</div></div>			
	F-W	Winter	High	<div><div></div><div>4%</div><div>13%</div></div>			
Yellow-bellied Sapsucker	F-E	Winter	Neutral	<div><div></div><div>4%</div><div></div></div>			
Red-naped Sapsucker	F-W	Summer	High	<div><div></div><div>35%</div><div>9%</div></div>			
	F-W	Winter	Neutral	<div><div></div><div>1%</div><div>31%</div></div>			
Lewis's Woodpecker	F-W	Summer	Moderate	<div><div></div><div>27%</div><div>19%</div></div>			
	F-W	Winter	Low	<div><div></div><div>9%</div><div>21%</div></div>			
Red-headed Woodpecker	F-E	Summer	High	<div><div></div><div>5%</div><div></div></div>			
Acorn Woodpecker	F-W	Summer	High	<div><div></div><div><1%</div><div>3%</div></div>			
	F-W	Winter	Moderate	<div><div></div><div>1%</div><div>2%</div></div>			
	F-B	Summer	High	<div><div></div><div>23%</div><div>4%</div></div>			





















































Species	Habitat Group	Season	Range-wide Vulnerability	State Trends	State Threats
American Three-toed Woodpecker	F-B	Winter	High	<div><div>24%</div><div>3%</div></div>	
Downy Woodpecker	Gen	Summer	Neutral	<div><div>31%</div><div>22%</div></div>	 
	Gen	Winter	Neutral	<div><div>30%</div><div>17%</div></div>	 
Ladder-backed Woodpecker	D	Summer	Neutral	<div><div><1%</div><div>32%</div></div>	 
	D	Winter	Neutral	<div><div><1%</div><div>41%</div></div>	 
Hairy Woodpecker	Gen	Summer	Low	<div><div>15%</div><div>18%</div></div>	 
	Gen	Winter	Low	<div><div>7%</div><div>17%</div></div>	 
Pileated Woodpecker	F-E	Summer	Neutral	<div><div>7%</div></div>	 
	F-E	Winter	Neutral	<div><div>4%</div></div>	 
Northern Flicker	Gen	Summer	Moderate	<div><div>43%</div><div>38%</div></div>	 
	Gen	Winter	Neutral	<div><div>63%</div><div>27%</div></div>	 
American Kestrel	Gen	Summer	Neutral	<div><div>84%</div><div>7%</div></div>	 
	Gen	Winter	Neutral	<div><div>49%</div><div>29%</div></div>	 
Merlin	F-E	Winter	Neutral	<div><div>76%</div><div>18%</div></div>	 
Aplomado Falcon	G	Winter	Moderate	<div><div>5%</div></div>	 
Peregrine Falcon	Gen	Summer	Neutral	<div><div>71%</div><div>18%</div></div>	 
	Gen	Winter	Neutral	<div><div>8%</div><div>37%</div></div>	 
Prairie Falcon	D	Summer	Low	<div><div>22%</div><div>7%</div></div>	 
	D	Winter	Low	<div><div>76%</div><div>4%</div></div>	 
Olive-sided Flycatcher	F-B	Summer	High	<div><div>22%</div><div>4%</div></div>	
Western Wood-Pewee	F-W	Summer	High	<div><div>28%</div><div>29%</div></div>	 
Willow Flycatcher	F-W	Summer	Moderate	<div><div>11%</div><div>14%</div></div>	 
Hammond's Flycatcher	F-W	Summer	High	<div><div>16%</div><div>5%</div></div>	 
Gray Flycatcher	D	Summer	High	<div><div>15%</div><div>14%</div></div>	 
Dusky Flycatcher	F-W	Summer	High	<div><div>21%</div><div>11%</div></div>	 
Cordilleran Flycatcher	F-W	Summer	High	<div><div>17%</div><div>15%</div></div>	 
Black Phoebe	Gen	Summer	Neutral	<div><div><1%</div><div>16%</div></div>	 

Species	Habitat Group	Season	Range-wide Vulnerability	State Trends	State Threats			
	Gen	Winter	Neutral	<div><div></div>4%</div>				
Say's Phoebe	Gen	Summer	Low	<div><div></div>60%<div></div>16%</div>				
Vermilion Flycatcher	D	Winter	Neutral	<div><div></div>5%</div>				
Ash-throated Flycatcher	D	Summer	Neutral	<div><div></div>9%<div></div>43%</div>				
Cassin's Kingbird	D	Summer	High	<div><div></div>3%<div></div>3%</div>				
Western Kingbird	G	Summer	Neutral	<div><div></div>44%<div></div>22%</div>				
Eastern Kingbird	G	Summer	Moderate	<div><div></div>8%</div>				
Scissor-tailed Flycatcher	G	Summer	Neutral	<div><div></div>2%<div></div>40%</div>				
Loggerhead Shrike	G	Summer	Neutral	<div><div></div>38%<div></div>17%</div>				
	G	Winter	Neutral	<div><div></div>6%<div></div>47%</div>				
Northern Shrike	F-B	Winter	Moderate	<div><div></div>61%<div></div>17%</div>				
Bell's Vireo	D	Summer	Low	<div><div></div><1%<div></div>35%</div>				
Gray Vireo	D	Summer	Moderate	<div><div></div>2%<div></div>15%</div>				
Cassin's Vireo	F-W	Summer	Low	<div><div></div>11%</div>				
Plumbeous Vireo	F-W	Summer	Neutral	<div><div></div>27%<div></div>21%</div>				
Warbling Vireo	Gen	Summer	Neutral	<div><div></div>34%<div></div>27%</div>				
Red-eyed Vireo	F-E	Summer	Low	<div><div></div>7%<div></div>8%</div>				
Canada Jay	F-B	Summer	High	<div><div></div>5%</div>				
	F-B	Winter	High	<div><div></div>13%<div></div>3%</div>				
Pinyon Jay	F-W	Summer	Moderate	<div><div></div>16%<div></div>13%</div>				
	F-W	Winter	Low	<div><div></div>15%<div></div>13%</div>				
Steller's Jay	F-W	Summer	Moderate	<div><div></div>8%<div></div>13%</div>				
	F-W	Winter	Moderate	<div><div></div>17%<div></div>8%</div>				
Blue Jay	F-E	Summer	Neutral	<div><div></div>1%<div></div>8%</div>				
Woodhouse's Scrub-Jay	F-W	Summer	Moderate	<div><div></div>19%<div></div>26%</div>				
	F-W	Winter	Moderate	<div><div></div>25%<div></div>31%</div>				
Mexican Jay	F-S	Summer	High	<div><div></div>4%</div>				
	F-S	Winter	Moderate	<div><div></div>2%</div>				
























































Species	Habitat Group	Season	Range-wide Vulnerability	State Trends		State Threats	
Black-billed Magpie	Gen	Summer	High	<div><div>32%</div><div>24%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
	Gen	Winter	Moderate	<div><div>40%</div><div>26%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Clark's Nutcracker	F-W	Summer	High	<div><div>27%</div><div>11%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
	F-W	Winter	High	<div><div>31%</div><div>7%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
American Crow	Gen	Summer	Low	<div><div>28%</div><div>27%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
	Gen	Winter	Neutral	<div><div>34%</div><div>37%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Chihuahuan Raven	D	Summer	Neutral	<div><div>14%</div><div>29%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
	D	Winter	Neutral	<div><div>3%</div><div>33%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Common Raven	Gen	Summer	Low	<div><div>15%</div><div>83%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
	Gen	Winter	Low	<div><div>29%</div><div>56%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Horned Lark	G	Summer	Low	<div><div>5%</div><div>54%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
	G	Winter	Low	<div><div>45%</div><div>6%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Northern Rough-winged Swallow	Gen	Summer	Neutral	<div><div>68%</div><div>14%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Purple Martin	Gen	Summer	Neutral	<div><div><1%</div><div>15%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Tree Swallow	Gen	Summer	Moderate	<div><div>25%</div><div>8%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Violet-green Swallow	F-W	Summer	Moderate	<div><div>10%</div><div>41%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Bank Swallow	Gen	Summer	Neutral	<div><div>15%</div><div>16%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Barn Swallow	Gen	Summer	Neutral	<div><div>57%</div><div>14%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Cliff Swallow	Gen	Summer	Neutral	<div><div>76%</div><div>18%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Black-capped Chickadee	F-B	Summer	Low	<div><div>50%</div><div>14%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
	F-B	Winter	Low	<div><div>50%</div><div>13%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Mountain Chickadee	F-W	Summer	High	<div><div>16%</div><div>24%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
	F-W	Winter	High	<div><div>20%</div><div>27%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Bridled Titmouse	F-S	Winter	High	<div><div>2%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div><div></div></div>	
Juniper Titmouse	F-W	Summer	Low	<div><div>2%</div><div>13%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
	F-W	Winter	Low	<div><div>4%</div><div>16%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div></div>	
Verdin	D	Winter	Neutral	<div><div>14%</div></div>	<div><div></div><div></div></div>	<div><div></div><div></div><div></div></div>	




















































Species	Habitat Group	Season	Range-wide Vulnerability	State Trends	State Threats
Bushtit	F-W	Summer	High	<div><div>7%</div><div>23%</div></div>	 
	F-W	Winter	Moderate	<div><div>4%</div><div>20%</div></div>	 
Red-breasted Nuthatch	F-B	Summer	Moderate	<div><div>23%</div><div>10%</div></div>	
	F-B	Winter	Neutral	<div><div>65%</div><div>16%</div></div>	 
White-breasted Nuthatch	F-E	Summer	Low	<div><div>42%</div><div>17%</div></div>	 
	F-E	Winter	Neutral	<div><div>43%</div><div>22%</div></div>	 
Pygmy Nuthatch	F-W	Summer	High	<div><div>20%</div><div>8%</div></div>	 
	F-W	Winter	Moderate	<div><div>14%</div><div>11%</div></div>	 
Brown Creeper	F-W	Summer	Moderate	<div><div>18%</div><div>10%</div></div>	 
	F-W	Winter	Neutral	<div><div>48%</div><div>39%</div></div>	 
Rock Wren	D	Summer	Moderate	<div><div>5%</div><div>34%</div></div>	 
	D	Winter	Neutral	<div><div>25%</div></div>	 
Canyon Wren	D	Summer	Neutral	<div><div>5%</div><div>20%</div></div>	 
	D	Winter	Neutral	<div><div>14%</div><div>31%</div></div>	 
House Wren	Gen	Summer	Moderate	<div><div>29%</div><div>18%</div></div>	 
Pacific Wren	F-W	Summer	Neutral	<div><div>2%</div></div>	  
Marsh Wren	M	Summer	Low	<div><div>2%</div><div>1%</div></div>	 
	M	Winter	Low	<div><div>4%</div><div>36%</div></div>	 
Bewick's Wren	D	Summer	Neutral	<div><div>13%</div><div>57%</div></div>	 
	D	Winter	Low	<div><div>9%</div><div>56%</div></div>	 
Cactus Wren	D	Summer	Neutral	<div><div>54%</div></div>	 
	D	Winter	Neutral	<div><div>16%</div></div>	  
Blue-gray Gnatcatcher	Gen	Summer	Neutral	<div><div>8%</div><div>20%</div></div>	 
Black-tailed Gnatcatcher	D	Summer	Neutral	<div><div>2%</div></div>	 
American Dipper	F-W	Summer	Moderate	<div><div>9%</div><div>39%</div></div>	 
	F-W	Winter	High	<div><div>24%</div><div>16%</div></div>	 
Golden-crowned Kinglet	F-B	Summer	Moderate	<div><div>8%</div><div>3%</div></div>	
	F-B	Winter	Neutral	<div><div>7%</div><div>8%</div></div>	 

































Species	Habitat Group	Season	Range-wide Vulnerability	State Trends		State Threats		
Ruby-crowned Kinglet	F-W	Summer	High	26% 7%				
	F-W	Winter	Neutral	50%				
Western Bluebird	F-W	Summer	Moderate	10% 21%				
	F-W	Winter	High	4% 11%				
Mountain Bluebird	F-W	Summer	High	16%				
	F-W	Winter	Low	33%				
Townsend's Solitaire	F-W	Summer	High	18% 11%				
	F-W	Winter	High	40% 35%				
Swainson's Thrush	F-B	Summer	High	<1% 4%				
Hermit Thrush	F-W	Summer	High	12% 8%				
	F-W	Winter	Low	<1% 23%				
American Robin	Gen	Summer	Moderate	47% 35%				
	Gen	Winter	Neutral	41% 29%				
Gray Catbird	F-E	Summer	Neutral	47% 11%				
Curve-billed Thrasher	D	Summer	Neutral	14% 35%				
	D	Winter	Neutral	11%				
Brown Thrasher	F-E	Summer	High	4%				
Bendire's Thrasher	D	Summer	Low	7%				
Crissal Thrasher	D	Summer	Low	26%				
	D	Winter	Low	17%				
Sage Thrasher	D	Summer	High	11% 7%				
	D	Winter	Low	13% 35%				
Northern Mockingbird	Gen	Summer	Neutral	20% 27%				
	Gen	Winter	Neutral	34%				
American Pipit	A	Summer	High	7%				
	A	Winter	Neutral	7% 26%				
Sprague's Pipit	G	Winter	Neutral	23%				
Bohemian Waxwing	F-B	Winter	High	32% 10%				

Species	Habitat Group	Season	Range-wide Vulnerability	State Trends		State Threats		
Cedar Waxwing	Gen	Summer	Low	24%	14%			
	Gen	Winter	Neutral	39%	20%			
Phainopepla	D	Summer	Neutral	7%				
	D	Winter	Neutral	8%				
Evening Grosbeak	F-B	Summer	High	20%	6%			
	F-B	Winter	Moderate	32%	9%			
Pine Grosbeak	F-B	Summer	High	7%	1%			
	F-B	Winter	Moderate	19%	3%			
Gray-crowned Rosy-Finch	A	Winter	High	38%	6%			
Black Rosy-Finch	A	Winter	High	35%	4%			
Brown-capped Rosy-Finch	A	Summer	High	3%				
	A	Winter	High	23%				
House Finch	Gen	Summer	Low	51%	29%			
	Gen	Winter	Low	49%	31%			
Purple Finch	F-B	Winter	Low	<1%	5%			
Cassin's Finch	F-W	Summer	High	29%	22%			
	F-W	Winter	Moderate	71%	9%			
Common Redpoll	A	Winter	Low	9%				
Red Crossbill	F-B	Summer	High	28%	9%			
	F-B	Winter	Moderate	38%	15%			
White-winged Crossbill	F-B	Summer	High	6%				
	F-B	Winter	Moderate	31%	5%			
Pine Siskin	F-W	Summer	Moderate	25%	11%			
	F-W	Winter	Neutral	16%	69%			
Lesser Goldfinch	F-W	Summer	Neutral	22%	41%			
	F-W	Winter	Neutral	7%	59%			
Lawrence's Goldfinch	D	Winter	Low	3%				
American Goldfinch	Gen	Summer	Moderate	1%				

Species	Habitat Group	Season	Range-wide Vulnerability	State Trends		State Threats	
	Gen	Winter	Neutral	54%	25%		
Lapland Longspur	A	Winter	Neutral	16%	20%		
Chestnut-collared Longspur	G	Summer	High	1%	9%		
	G	Winter	Moderate	37%			
Smith's Longspur	A	Winter	Neutral	16%			
McCown's Longspur	G	Summer	High	4%			
	G	Winter	Moderate	19%			
Snow Bunting	A	Winter	Low	51%	14%		
Cassin's Sparrow	G	Summer	Low	34%	9%		
Grasshopper Sparrow	G	Summer	Low	26%	8%		
	G	Winter	Neutral	13%			 
Chipping Sparrow	Gen	Summer	Moderate	15%	16%		
	Gen	Winter	Neutral	5%			 
Clay-colored Sparrow	G	Winter	Neutral	2%			 
Black-chinned Sparrow	D	Summer	High	2%			
	D	Winter	Low	21%			
Brewer's Sparrow	D	Summer	High	23%	9%		
	D	Winter	Moderate	5%			 
Black-throated Sparrow	D	Summer	Neutral	8%	44%		
	D	Winter	Neutral	9%	49%		
Lark Sparrow	D	Summer	Neutral	53%	20%		
Lark Bunting	G	Summer	High	34%	13%		
	G	Winter	Neutral	29%			
American Tree Sparrow	A	Winter	Neutral	24%	39%		
Fox Sparrow	F-B	Summer	High	4%			
Dark-eyed Junco	F-W	Summer	High	24%	11%		
	F-W	Winter	Neutral	59%	31%		
White-crowned Sparrow	Gen	Summer	High	25%	2%		

Species	Habitat Group	Season	Range-wide Vulnerability	State Trends		State Threats	
	Gen	Winter	Neutral	28%	32%		
Harris's Sparrow	F-B	Winter	Low	4%	11%		
White-throated Sparrow	F-B	Winter	Neutral	8%	40%		
Sagebrush Sparrow	D	Summer	High	7%	6%		
	D	Winter	Neutral	18%			
Vesper Sparrow	G	Summer	Moderate	22%	15%		
	G	Winter	Neutral	35%			
Savannah Sparrow	G	Summer	High	6%	<1%		
	G	Winter	Low	28%			
Song Sparrow	Gen	Summer	Moderate	42%	16%		
	Gen	Winter	Neutral	39%	27%		
Lincoln's Sparrow	F-B	Summer	High	24%	3%		
	F-B	Winter	Neutral	20%	50%		
Swamp Sparrow	M	Winter	Neutral	<1%	2%		 
Canyon Towhee	D	Summer	Low	8%	30%		
	D	Winter	High	<1%	5%		
Rufous-crowned Sparrow	D	Summer	Low	<1%	13%		
	D	Winter	High	<1%	5%		
Green-tailed Towhee	D	Summer	High	23%	10%		
	D	Winter	Neutral	2%			 
Spotted Towhee	F-W	Summer	Moderate	13%	19%		
	F-W	Winter	Low	11%	45%		
Yellow-breasted Chat	F-E	Summer	Neutral	17%	42%		
Yellow-headed Blackbird	M	Summer	Low	9%	9%		
	M	Winter	Low	5%			 
Western Meadowlark	G	Summer	Low	11%	47%		
	G	Winter	Neutral	24%	19%		
Eastern Meadowlark	G	Summer	Moderate	17%			

Species	Habitat Group	Season	Range-wide Vulnerability	State Trends	State Threats		
	G	Winter	Neutral	25%			
Orchard Oriole	F-E	Summer	Low	23% 1%			
Hooded Oriole	F-W	Summer	Neutral	2%			
Bullock's Oriole	F-W	Summer	Neutral	46% 24%			
Baltimore Oriole	F-E	Summer	Low	15% 2%			
Scott's Oriole	D	Summer	Neutral	3% 26%			
Red-winged Blackbird	Gen	Summer	Neutral	54% 7%			
	Gen	Winter	Neutral	49% 26%			
Brown-headed Cowbird	Gen	Summer	Neutral	73% 11%			
	Gen	Winter	Neutral	22%			
Brewer's Blackbird	Gen	Summer	Moderate	39% 35%			
	Gen	Winter	Neutral	17% 58%			
Common Grackle	F-E	Summer	Low	33% 2%			
Great-tailed Grackle	Gen	Summer	Neutral	12% 35%			
	Gen	Winter	Neutral	2% 34%			
Northern Waterthrush	F-B	Summer	Moderate	3%			
Orange-crowned Warbler	F-W	Summer	High	10% 6%			
	F-W	Winter	Neutral	2%			
Lucy's Warbler	D	Summer	Low	14%			
Virginia's Warbler	F-W	Summer	Moderate	7% 17%			
MacGillivray's Warbler	F-W	Summer	Moderate	12% 9%			
Common Yellowthroat	Gen	Summer	Low	17% 13%			
American Redstart	F-B	Summer	Moderate	10% 7%			
Yellow Warbler	F-B	Summer	Moderate	33% 13%			
Pine Warbler	F-E	Summer	High	2%			
Yellow-rumped Warbler	F-B	Summer	Moderate	23% 9%			
	F-B	Winter	Neutral	31% 44%			

Species	Habitat Group	Season	Range-wide Vulnerability	State Trends	State Threats
Black-throated Gray Warbler	F-W	Summer	Moderate	<div><div>10%</div><div>19%</div></div>	 
Wilson's Warbler	F-W	Summer	High	<div><div>12%</div><div>3%</div></div>	 
Red-faced Warbler	F-S	Summer	High	<div><div>4%</div></div>	 
Hepatic Tanager	F-W	Summer	Moderate	<div><div>2%</div><div>6%</div></div>	  
Summer Tanager	F-E	Summer	Neutral	<div><div>6%</div></div>	  
Western Tanager	F-W	Summer	Moderate	<div><div>10%</div><div>24%</div></div>	 
Northern Cardinal	F-E	Summer	Neutral	<div><div>33%</div><div>8%</div></div>	 
	F-E	Winter	Neutral	<div><div><1%</div><div>16%</div></div>	 
Pyrrhuloxia	D	Summer	Neutral	<div><div>2%</div></div>	  
	D	Winter	Neutral	<div><div>3%</div></div>	  
Black-headed Grosbeak	F-W	Summer	Moderate	<div><div>10%</div><div>22%</div></div>	 
Blue Grosbeak	F-S	Summer	Neutral	<div><div>19%</div><div>33%</div></div>	 
Lazuli Bunting	F-W	Summer	Neutral	<div><div>25%</div><div>12%</div></div>	 
Painted Bunting	D	Summer	Neutral	<div><div>10%</div></div>	 
Dickcissel	G	Summer	Neutral	<div><div>10%</div><div>12%</div></div>	